

# How to Interpret $Z_{DR}$ Shade Charts


Lindsey M. Richardson, W. David Zittel, Dan  
B. Frashier, Amy E. Daniel, and Jessica A.  
Schultz

Build 22+



# What's a Shade Chart?

- A graphical way to monitor  $Z_{DR}$  bias from a **single** radar site
  - Information from the most recent 6 months



# What's a Shade Chart? (Cont.)

- Based on 3 independent external target methods:
  - Light Rain
  - Dry Snow
  - Bragg Scatter
- Event characteristics are different between the methods



# Why do we care about $Z_{DR}$ Bias\*?

- $Z_{DR}$  bias shows the amount of error in ZDR Offset
- $Z_{DR}$  bias can have adverse affects on Quantitative Precipitation Estimation (QPE)
  - A positive  $Z_{DR}$  bias results in underestimation
  - A negative  $Z_{DR}$  bias results in overestimation
- $Z_{DR}$  affects other products as well
  - Melting Layer Detection Algorithm (MLDA)
    - Particularly “wet snow”
  - Hydrometeor Classification Algorithm (HCA)
    - Specific  $Z_{DR}$  thresholds for categories

\* This is not the same as ZDR Offset but is related





# Why external targets?

- External targets act as an estimation metric independent from the built-in hardware estimates
  - Additional measurement to verify built-in hardware results
- Methods work with operational scanning strategies and products



# What's on a Shade Chart?

Let's focus on one of the methods for more details




# Did you notice?

- The dashed lines?
- The black numbers at the top of the shading in places?
- The gap in shading?
- Each subplot contains information from an independent method?
- The chart has a trend in time?


Let's focus on a single subplot and explore these details

# Layout of Each Subplot






# $Z_{DR}$ Bias Estimates from Events



- Many, highly-varying scatter points
  - Events may vary greatly from one to another
  - Events are defined per method in later slides
- Events are OK, but what if we took a 7-day median?


# Grouping for Shading

Site A Systematic  $Z_{DR}$  Bias Jun'15-Nov'15  
Based on  $\Delta_{Rain}$  (dB)




Zoom in to focus on smaller time scale

# 7-Day Running Median




- For each 7-day grouping, a median is calculated from the points
  - Days 1-7, 2-8, 3-9, etc.
- Shading vertex placed on middle day of 7-day set
- Each of the colored boxes above represent a separate set

# 7-Day Median Calculation




Go from just having points...

# 7-Day Median Calculation




...to having shading.

# 7-Day Median Shading



- Long-term median trend shows a continuing bias
  - Positive (red-shaded) values are considered warm or high
  - Negative (blue-shaded) values are considered cool or low
- Sites are beyond recommended limits if the shading is regularly above (below) the dashed 0.2 dB (-0.2 dB) line


# 7-Day Median Shading Cont.



- Medians beyond the y-axis limits are shown as a number near the top
- No shading means the data is either missing, not available, or equals 0.0 dB exactly
- Will interpolate if only missing one shade value between two valid points

# 7-Day Median Shading Cont.

Where are the scatter points?




- Event scatter points outside of the y-axis limits ( $\pm 0.5$  dB) are not shown
- Recall that **median** values from shading outside of the limits are represented by the black numbers at the top




# Compare Trends from Multiple External Target Methods


Light Rain



Dry Snow




Bragg




# What do the methods detect?

Each method is independent and includes distinct caveats

- 2 Methods Related to Precipitation:
  - Light Rain (liquid precipitation)
  - Dry Snow (frozen precipitation)



- 1 Method Related to Clear Air
  - Bragg scattering associated with refractivity gradients



Click [here](#) to skip method details and jump to interpretation

# Light Rain Method

- Light rain has characteristics of slightly-wide (slightly-oblate) spheres
- Base Filters:
  - $\text{SNR} \geq 20 \text{ dB}$
  - $19.0 < Z < 21.0$
  - $0.98 < \rho_{\text{HV}} < 1.00$
  - Range Gate must at least 1 km below the bottom of the Melting Layer
- Additional Filters
  - Total Range Gates that pass Base Filters  $\geq 600$
  - $0.125 \leq \text{ZDR}_{\text{IQR}} \leq 0.35 \text{ dB}$
  - $0.0625 \leq \text{ZDR}_{\text{MEDAD}} \leq 0.25$
  - $15.0 \leq \text{Z90th} \leq 23.0 \text{ dBZ}$
  - $8.0 \leq \text{Z}_{\text{IQR}} \leq 16.0 \text{ dB}$
  - $0.3 \leq \text{PHI}_{\text{IQR}} \leq 3.3^\circ$



# Light Rain Method Cont.

- Daily Median (scatter points on chart)
  - An estimate is calculated each volume via:  
ZDR Mode – ZDR Intrinsic
    - ZDR Intrinsic for Light Rain in this reflectivity range is **0.25 dB**
  - The most recent 12 volumes are averaged together
    - Both the single volume and 12-volume average can be found in the RPG Status Log or in the ASP on lines that begin with “ZDRBE”
  - For a single date (based on UTC time), the median of the 12-volume averages defines the Daily Median
- Convective processes and other particle contamination can **bias  $Z_{DR}$  estimates high**
- Intrinsic ZDR Subtraction factor can **bias  $Z_{DR}$  estimates low**

\* The ASP is a product version of the RPG Status Log available from archive sources such as NCEI.

# Dry Snow Method


- Only uses range gates classified as Dry Snow (dry aggregates) by the Hydrometeor Classification Algorithm (HCA)
  - Dry aggregates appear similar to spheroids when they fall
  - Snow does NOT have to be reaching the surface
- Base Filters:
  - Range > 20 km
  - $15 \text{ dBZ} < Z < 25 \text{ dBZ}$
  - Elevations >  $1^\circ$
  - SNR  $\geq 20 \text{ dB}$ ,
  - $0.98 < \text{RHO}_{\text{HV}} < 1.0$
  - $\text{PHI} < 100^\circ$
  - Range gate must be within the first 1km completely above the melting layer
- Additional Filters
  - Must have at least 600  $Z_{\text{DR}}$  range gates that pass filters per volume
  - Standard deviation  $Z_{\text{DR}} < 0.5 \text{ dB}$

Note: Aggregates are clumps of frozen precipitation (particularly ice crystals)




# Dry Snow Method Cont.

- Daily Median Events
  - An estimate is calculated each volume via:  
ZDR Mode – ZDR Intrinsic
    - ZDR Intrinsic used for Dry Aggregate Snow is **0.20 dB**
  - The most recent 12 volumes are averaged together
    - Both the single volume and 12-volume average can be found in the RPG Status Log or in the ASP on lines that begin with “ZDRBE”
  - For a single date (based on UTC time), the median of the 12-volume averages defines the Daily Median
- Can be estimated at the same time as a rain Event as long as dry aggregate snow is observed above the melting layer
- Dendrites and Platelets can **bias  $Z_{DR}$  estimates high**
- Subtraction factor can **bias  $Z_{DR}$  estimates low**



# Bragg Scatter Method

- Bragg distinguished by refractivity gradients generally caused by turbulent eddies
  - Often found at the top of the Convective Boundary Layer and Marine Boundary Layer
- Base Filters:
  - 10-80 km in range only
  - $Z < 10$  dBZ
  - $|V| > 2$  m/s
  - $W > 0$  m/s
  - $SNR < 15$  dB
  - $0.98 < \rho_{HV} < 1.05$
  - Elevations  $2.4-4.5^\circ$
- Additional Filters:
  - Must have at least 600  $Z_{DR}$  range gates that pass filters per volume
  - $Z$  at the 90<sup>th</sup> percentile  $\leq -3$  dBZ (precipitation filter)
  - Inter-Quartile Range (IQR)  $< 0.9$  (biota filter)



# Bragg Scatter Method Cont.

- Daily Median Events
  - An estimate is calculated each volume via:  
ZDR Mode – ZDR Intrinsic
    - ZDR Intrinsic used for Bragg Scatter is **0.00 dB**
  - The most recent 12 volumes are averaged together but only if there are at least 10,000 range gates of potential Bragg estimates in those 12 volumes.
    - Both the single volume and 12-volume average can be found in the RPG Status Log or in the ASP on lines that begin with “ZDRBE”
  - For a single date (based on UTC time), the median of the 12-volume averages defines the Daily Median
- Precipitation contamination can **bias  $Z_{DR}$  high**
- Return from Bragg scattering has a weak signal, and if noise is comparable to the signal it could **bias the estimate towards 0.0 dB**
  - Assuming the noise estimates are similar in both H and V channels



# Method Availability

- Light rain is less available during the cool season, especially at northern continental sites
- Dry snow can be found year round at most sites
- Bragg scatter less available due to stringent filters
  - Also less available in the warm season due to biota (bugs, birds, etc.) contamination




# Rely on More Than One Method When Possible!

- When all 3 methods show a similar bias, there is high confidence in the indicated bias
  - All methods are not always available
- The **trend** is the important aspect
  - Recommend **at least two weeks** of data to establish a baseline



# Full-Chart Recap: Compare the Methods

- $Z_{DR}$  is high (above and near the positive limit) for the majority of the time in all three methods
- Focus on most recent month
- Compare with previous months for overall trend



# Monthly Summary


- The median of each method is calculated for the most recent month (not shown)
- These medians are then used to calculate a Weighted Mean (WMean) Estimate (displayed in the top-right box)
  - Weights are based on method estimation accuracy
    - Bragg scatter: 42% weight
    - Dry Snow: 33% weight
    - Light Rain: 25% weight

Note: Volume-based Weighted Means using the same weights are shown on ZDRBE Status lines.



# Monthly Summary


- The monthly WMean box is color coded (matches shade color when beyond-recommended-limits)
- It will say NaN (Not-a-Number) if there are no estimates for an entire month



# Shade Interpretation: “Good” Site

Within Limits


Close to zero  
and within  
recommended  
limits for all  
methods



# Shade Interpretation: Another “Good” Site

Within Limits


Alternating  
red/blue  
around zero  
(within  
limits) is OK



# Shade Interpretation: Warm Bias

## Beyond Limits

A warm bias  
causes QPE  
underestimation






# Shade Interpretation: Cold Bias


Beyond Limits

A cold bias  
causes QPE  
overestimation



# Shade Interpretation: Site Improves


Site had a large negative bias and improved closer to within limits



# Shade Interpretation: Disagreement

Disagreement possible due to the independent method caveats

- Site bias is likely around 0.0 dB in this case
- Within limits on both sides, so less priority to take action




# 6-Month Time Window

- Shading features are consistent for a given month

Site H  
End Month:  
September


Note how  
the circled  
areas move  
with time in  
the next two  
slides



# 6-Month Time Window

- Features move to the left


Site H  
End Month:  
October



# 6-Month Time Window

- Some features move off with the moving time window

Site H  
End Month:  
November




Previous  
months are  
archived and  
available




# Redundant Sites (Ch1 Example)

Attempt to isolate estimates per channel  
- The ROC will send charts for each channel



# Redundant Sites (Ch2 example)

Sometimes the plots only have info in one Channel. This may be a plotting error if the site is routinely switching channels as suggested.






# Shade Chart Availability

- Shade Charts are available with frequent updates via the Data Quality Dashboard on the MSCF
  - Click on the big Main Gauge or select from the menu
- They are also available from the ROC webpage and updated monthly:
  - <https://www.roc.noaa.gov/branches/operations-branch/wsr88d-hotline-links-info.php>

# Shade Chart in DQD




# Summary

- A shade chart is a quick way to assess if a site has a  $Z_{DR}$  bias and the approximate magnitude of the bias
  - Sites with biases outside of the  $\pm 0.2$  dB range are considered to be **Beyond Recommended Limits**
  - $Z_{DR}$  bias adversely affects several products, especially QPE
- Charts can help track when maintenance was performed and if it helped (e.g., had a large bias and was corrected to within limits)
  - Can also see if a site has a new or drifting hardware issue (e.g., site was within limits and jumped to a large bias)



# Summary Cont.

- We are still exploring the details of the external target methods!
  - External targets are an independent, extra metric to the built-in hardware estimates that work with operational scanning strategies and products
  - Each method has unique caveats and variability in accuracy
    - Some aspects of variability remain unknown
- By using multiple methods, there is higher confidence a site does or does not have a bias



# Within Limits is OK!

- Achieving an exact  $Z_{DR}$  bias estimates of 0.0 dB can be difficult because the variability of the methods and built-in hardware often exceed  $\pm 0.1$  dB
  - The trend of median bias estimates falling within  $\pm 0.2$  dB should be sufficient for most algorithms and visual analysis



# Extra Information

- Read publications and more by visiting the WSR-88D Hotline site:

<https://www.roc.noaa.gov/branches/operations-branch/wsr88d-hotline-links-info.php>

- Other articles can be found at ROC Papers:

<https://www.roc.noaa.gov/branches/field-requirements-branch/reports-and-conferences.php>

